

Mesoscopic charge fluctuations in the Coulomb blockade regime

A. Kaminski¹, I.L. Aleiner², and L.I. Glazman¹

¹*Theoretical Physics Institute, University of Minnesota, Minneapolis, MN 55455*

²*Department of Physics, State University of New York, Stony Brook, NY 11794*

We study mesoscopic fluctuations of the differential capacitance of a dot connected to a lead by a single-mode quantum channel with adjustable conductance G . We show that the amplitude of the fluctuations reaches maximum at a partially opened channel, $G \lesssim e^2/\pi\hbar$. Parametric correlations of fluctuations at different values of the applied voltage and magnetic field can be studied experimentally, and we find the corresponding correlation functions for the differential capacitance.

PACS numbers: 73.23.-b, 73.23.Hk, 73.40.Gk

Charge of a conducting dot weakly connected to leads is quantized in the units of electron charge e . When the potential induced on the dot is varied, its charge varies in a step-like manner. (The potential can be tuned by means of voltage V_g applied to a gate, which is coupled to the dot only electrostatically.) The step-like behavior of the dot charge $Q(V_g)$, *i.e.*, Coulomb blockade, was clearly demonstrated in experiments utilizing metallic islands [1]. The charge quantization is progressively smeared with the increase of the conductance G_0 of a junction connecting dot with a lead. Theory [2] predicts that deviations of Q from the quantized values are proportional to conductance, $|Q(V_g) - me| \propto G_0$ at $G_0 \ll G_q$. When G_0 is of the order of the quantum unit $G_q \equiv e^2/\pi\hbar$, the function $Q(V_g)$ depends on the detailed properties of the junction [3–5]. For a single-mode junction, the Coulomb blockade vanishes when the reflectionless propagation of an electron mode through the dot–lead channel is almost reached, with the periodic part of $Q(V_g)$ approaching zero approximately as $G_q - G_0$, see [5]. Single-mode junction allowing for ballistic propagation of electrons is a reasonable model for a semiconductor quantum dot device. In such a device, G_0 is tunable, unlike in the lithographically prepared metallic samples [1]. This allowed several groups to attack the challenging problem of quantum charge fluctuations in the Coulomb blockade regime. Suppression of the Coulomb blockade effect at $G_0 \rightarrow G_q$ was demonstrated experimentally by Waugh *et al.* [6] and Molenkamp *et al.* [7] on double-dot devices. In very recent experiments, Berman *et al.* [8] measured the charge smearing in a single dot.

Mesoscopic fluctuations of the charge Q and correspondingly of the differential capacitance C_{diff} of a dot reflect the randomness of the electron states within the dot. Gopar *et al.* [9] employed the Random Phase Approximation to consider the charge dynamics of the dot connected to a reservoir by a reflectionless ($G_0 = G_q$) one mode channel. In this approach, the mesoscopic fluctuations are associated with the fluctuations of the density of states on the Fermi level. Later it was shown [10] that even in this regime the Coulomb interaction leads to the formation of a collective state involving one-

electron states in a wide energy strip around the Fermi level (width of the strip is of the order of the charging energy E_C). As the result, the mesoscopic fluctuations of the capacitance at $G_0 = G_q$ are larger than predicted in Ref. [9], and the scale of the correlation magnetic field is controlled by the charging energy rather than by one electron level spacing.

In this Letter, we will study the capacitance fluctuations in the whole region $0 < G_0 < G_q$. We will show that the mesoscopic fluctuations are determined by the fluctuations of a large number ($\simeq E_C/\Delta$, where Δ is the one-electron level spacing) of single electron wave functions rather than by the state at the Fermi level. Concentrating on the low temperature limit $T \lesssim \Delta$, we show that the dependence of the variance $\langle \delta C_{diff}^2 \rangle$ on conductance G_0 is non-monotonous, with the maximum of fluctuations reached when the channel is partially open, $G_0 < G_q$. We will find explicit analytic expressions for the correlation functions characterizing the evolution of the fluctuating differential capacitance with the magnetic field.

We account for the Coulomb interaction by adding an extra term to the Hamiltonian:

$$\hat{H} = \hat{H}_0 + E_C(\hat{n} - \mathcal{N})^2. \quad (1)$$

Here $\hat{H}_0$ is the Hamiltonian of the dot-lead system without Coulomb interaction, and the term $E_C(\hat{n} - \mathcal{N})^2$ accounts for the charging energy of the dot, $\hat{n} = \sum_p c_p^\dagger c_p$ is the total number of electrons in the dot. The gate voltage V_g is represented by the conventional dimensionless parameter $\mathcal{N} = V_g C_g / e$, with C_g being the gate capacitance. The differential capacitance of the dot C_{diff} is related to the ground state energy E_0 of the Hamiltonian (1) by

$$C_{diff}(\mathcal{N}) \equiv \frac{C}{C_g} \frac{\partial Q}{\partial V_g} = C \left(1 - \frac{1}{2E_C} \frac{\partial^2 E_0}{\partial \mathcal{N}^2} \right). \quad (2)$$

Here Q is the dot charge, $e\hat{n}$, averaged over the ground state of the system, C is the capacitance of the isolated dot; we have introduced a lever-arm coefficient C/C_g into the definition of C_{diff} .

The strength of tunneling through the dot-lead junction is characterized by its dimensionless conductance

$g \equiv G_0/G_q$. To demonstrate the nonmonotonous behavior of the amplitude of the differential capacitance fluctuations, we consider analytically the limits of weak tunneling ($g \ll 1$) and of almost ballistic propagation of electrons in the junction ($1 - g \ll 1$).

Weak tunneling. The case of the weak tunneling, $g \ll 1$, is adequately described in the tunneling Hamiltonian formalism,

$$\begin{aligned} \hat{H}_0 = & \sum_k \xi_k a_k^\dagger a_k + \sum_p \xi_p c_p^\dagger c_p \\ & + \sum_{k,p} (t_{kp} a_k^\dagger c_p + t_{kp}^* c_p^\dagger a_k), \end{aligned} \quad (3)$$

where ξ_k and ξ_p are the one-electron energy spectra of the lead and dot respectively. Tunneling matrix elements here are related to g by $g = \nu_l \nu_d \langle |t_{kp}|^2 \rangle / 2\pi^2$. Hereafter $\langle \dots \rangle$ denotes the ensemble averaging, and ν_l and ν_d are the averaged densities of states in the lead and dot respectively. In the extreme $g \rightarrow 0$, the charge of the dot in the ground state is integer, $Q = me$. The half-integer values of the dimensionless voltage $\mathcal{N}$ correspond to the degeneracy points of the charging energy in Eq. (1). If V_g is tuned away from a degeneracy point ($m - 1/2 < \mathcal{N} < m + 1/2$), it takes a finite energy $2E_C(\mathcal{N} - m \pm 1/2)$ to increase or decrease the number of electrons in the dot by one with respect to the ground state.

The perturbation theory in the tunneling Hamiltonian (3) can be used to calculate the dot capacitance (2), if the system is not too close to one of the charge degeneracy points [2]:

$$g^{\frac{1}{2}} \ln(1/\mathcal{U}) \ll 1. \quad (4)$$

Here $\mathcal{U} \equiv \min\{|\mathcal{N} - m \pm 1/2|\}$. The lowest-order correction to the ground state energy is

$$E_{osc}(\mathcal{N}) = \frac{i}{2\pi} \frac{g}{\nu_{l0}\nu_{d0}} \int_{-\infty}^{\infty} dt \mathcal{G}_d(t; \mathcal{N}) \mathcal{G}_l(-t) \Big|_{\mathbf{r}=\mathbf{R}_c}, \quad (5)$$

where $\mathcal{G}_d$ and $\mathcal{G}_l$ are the one-electron Green functions in the dot and the lead, determined by the Hamiltonians $\hat{H}_d$ and $\hat{H}_l$ respectively, and $\mathbf{R}_c$ is the point of the dot-lead contact. The density of states per unit area in the dot and in the lead are denoted here as ν_{d0} and ν_{l0} . The dependence of $\mathcal{G}_d$ on the gate voltage is given by

$$\begin{aligned} \mathcal{G}_d(t, \mathcal{N}) = & \mathcal{G}_{d0}(t) \left[\theta(t) e^{-2i(m+1/2-\mathcal{N})E_C t} \right. \\ & \left. + \theta(-t) e^{2i(\mathcal{N}-m+1/2)E_C t} \right], \end{aligned} \quad (6)$$

where $\mathcal{G}_{d0}(t)$ is the one-electron Green function in the dot determined by the Hamiltonian $\hat{H}_{d0} = \sum_p \xi_p c_p^\dagger c_p$. The exponential factors account for the change in the Coulomb energy of the system when an extra electron or hole tunnels into the dot.

It follows from Eqs. (2), (5), and (6) that the fluctuations of the differential capacitance δC_{diff} are caused by the fluctuations of the one-electron Green function $\delta \mathcal{G}_{d0} \equiv \mathcal{G}_{d0} - \langle \mathcal{G}_{d0} \rangle$ in the dot at the point of the contact. Then the statistical properties of δC_{diff} can be related to the correlation function $\langle \delta \mathcal{G}_d(t_1; B_1) \delta \mathcal{G}_d(t_2; B_2) \rangle|_{\mathbf{R}_c}$. The applied magnetic field B affects the Green function $\mathcal{G}_{d0}$ thus affecting the capacitance fluctuations.

For a conducting dot, the correlation function for $\delta \mathcal{G}_{d0}$ is conventionally expressed in terms of the classical probabilities, diffuson $\mathcal{D}$ and Cooperon $\mathcal{C}$ [11]:

$$\begin{aligned} \langle \delta \mathcal{G}_{d0}(\omega_1; B_1) \delta \mathcal{G}_{d0}(\omega_2; B_2) \rangle|_{\mathbf{R}_c} = & 2\pi \nu_{d0} \theta(-\omega_1 \omega_2) \\ & \times [\mathcal{D}^{B_1, B_2}(|\omega_1 - \omega_2|) + \mathcal{C}^{B_1, B_2}(|\omega_1 - \omega_2|)]|_{\mathbf{R}_c}. \end{aligned} \quad (7)$$

For times exceeding the time of electron propagation across the dot of area S or, equivalently, at energies much less than the Thouless energy for a closed dot, E_T , the probability density of an electron is homogeneously distributed over the dot, and

$$\mathcal{D}^{B_1, B_2}(\omega) = \frac{S^{-1}}{-i\omega + \Omega_-}, \quad \mathcal{C}^{B_1, B_2}(t) = \frac{S^{-1}}{-i\omega + \Omega_+}. \quad (8)$$

Here $\Omega_{\pm} = E_T (S(B_1 \pm B_2)/\Phi_0)^2$, and $\Phi_0 = 2\pi\hbar c/e$ is the flux quantum. Differentiating Eq. (5) twice over $\mathcal{N}$ and extracting the random part, we come to the expression that relates fluctuations of the dot differential capacitance δC_{diff} to the fluctuations of the one-electron Green function in the dot $\delta \mathcal{G}_d$. The correlation functions for δC_{diff} are thus related to those for $\delta \mathcal{G}_d$, which are given by Eqs. (6)–(8).

First we consider the variation of the dot capacitance with the magnetic field at the same gate voltage. The corresponding correlation function is

$$\begin{aligned} \langle \delta C_{diff}(\mathcal{N}, B_1) \delta C_{diff}(\mathcal{N}, B_2) \rangle \\ = \frac{C^2 g^2 \Delta}{24\pi^4 E_C \mathcal{U}^3} \sum_{\pm} M\left(\frac{B_{\pm}}{B_c(\mathcal{U})}\right) \end{aligned} \quad (9)$$

with $B_{\pm} = |B_1 \pm B_2|$, $\mathcal{U} \equiv \min\{|\mathcal{N} - m \pm 1/2|\}$, and

$$M(x) = \frac{6}{\pi} \int_0^{\infty} dy \frac{x^2}{y^2 + x^4} \frac{1}{y^3} \left[\frac{y^2 + 2y}{y + 1} - 2 \ln(1 + y) \right].$$

The function $M(x) \propto 1/x^2$ at large values of the argument $x \gg 1$, so that the correlation function decays with the increase of $B_{\pm}$ over the value B_c . At zero magnetic field, the variance of the capacitance is

$$\sqrt{\langle \delta C_{diff}^2 \rangle} = \frac{1}{2\sqrt{3}\pi^2} C g \sqrt{\frac{\Delta}{E_C}} \frac{1}{\mathcal{U}^{\frac{3}{2}}}.$$

The characteristic scale for the variation of the correlation function Eq. (9) with the magnetic field is [12]

$$B_c = \frac{\Phi_0}{S} \sqrt{\frac{U}{E_T}}. \quad (10)$$

The meaning of this formula for the correlation field B_c can be clarified if we recall the relation of the charge fluctuations to the electron Green function. In the semiclassical approximation, $\mathcal{G}(t) \propto e^{i\mathcal{S}(t)}$, where $\mathcal{S}$ is the action calculated along the trajectory. The dwelling time of an extra electron in the dot is $t_d \sim \hbar/U$. Over this time, the trajectory of a diffusing electron would cover an area $Dt_d \sim \hbar D/U$, if there were no bounds imposed by the dot size. The ratio $n_d = \hbar D/SU \simeq E_T/U$ determines the typical number of “winds” for the electron trajectory inside the dot. Because of the random direction of the winds, the total flux threading the electron trajectory is $\Phi = BS\sqrt{n_d}$, and the correction to the action due to magnetic field is $\mathcal{S}_B \cong 2\pi\Phi/\Phi_0$. Equating $\mathcal{S}_B$ to 2π , we obtain the formula for B_c .

Relatively short excitation lifetime, $t_d \ll \hbar/\Delta$, leads to correlations of the capacitance fluctuations in different Coulomb blockade valleys. In a valley, $\sim \hbar/\Delta t_d$ levels of spatial quantization closest to the Fermi level in the dot contribute to the charge fluctuations. Each subsequent peak of the Coulomb blockade corresponds to one more level occupied. Therefore, the fluctuations of capacitance at different valleys are correlated, unless the distance n between them exceeds E_C/Δ , so that an entirely new set of discrete levels contributes to the fluctuations. Calculation of the corresponding correlation function yields:

$$\langle \delta C_{diff}(\mathcal{U}) \delta C_{diff}(\mathcal{U} + n) \rangle \sim \frac{C^2 g^2 \Delta}{4\pi^4 E_C \mathcal{U}^3} \frac{\ln \left(1 + \frac{n\Delta}{2E_C \mathcal{U}} \right)}{\frac{n\Delta}{E_C \mathcal{U}}}. \quad (11)$$

It is to be mentioned that Eqs. (5) and (9) are valid only if condition (4) is satisfied. If $\mathcal{N}$ is close enough to one of the charge degeneracy points, so that (4) is violated (but $\mathcal{U}$ is still much greater than Δ/E_C), the finite-order perturbation theory is insufficient for calculation of the dot capacitance. Utilizing the renormalization group technique [2], we found that the capacitance fluctuations in this case are described by the same Eq. (9) with the renormalized junction conductance

$$\tilde{g} = g \left[\cos^2 \left(\frac{\sqrt{2g}}{\pi} \ln \frac{1}{\mathcal{U}} \right) \right]^{-1}. \quad (12)$$

With this modification, the condition for the applicability of Eq. (9) is relaxed from (4) to a weaker one, $g^{\frac{1}{2}} \ln(1/\mathcal{U}) \lesssim 1$. This new condition is imposed by the requirement that the renormalized conductance $\tilde{g}$ must be still much less than unity.

Strong tunneling. As it was pointed out in the introduction, the effects of the Coulomb blockade become less pronounced as the dimensionless conductance of the dot-lead junction, g , approaches unity. In the limit $\Delta = 0$, Flensberg [4] and Matveev [5] studied the case of a dot

connected to the lead by a constriction with small reflection amplitude $r = \sqrt{1-g}$. Since the characteristic energy E_C is much smaller than the Fermi energy, one can linearize the spectrum of one-dimensional electrons in the constriction. The electrons in the constriction are thus divided into left- and right-moving species, $\psi(x) = e^{-ik_F x} \psi_L(x) + e^{ik_F x} \psi_R(x)$. In [5], the full Hamiltonian (1) describing the electrons in the dot and lead connected to each other, is transformed to an effective one-dimensional form and expressed in terms ψ_L, ψ_R . In the limit $\Delta \rightarrow 0$ and in the absence of backscattering in the constriction, there is no mixing between the L and R states:

$$\hat{H}_{1D} = iv_F \int_{-\infty}^{\infty} dx \left\{ \psi_L^\dagger \partial_x \psi_L - \psi_R^\dagger \partial_x \psi_R \right\} + E_C \left(\int_{-\infty}^0 dx : \psi_L^\dagger \psi_L + \psi_R^\dagger \psi_R : - \mathcal{N} \right)^2; \quad (13)$$

here v_F is the Fermi velocity, and $: \dots :$ stands for the normal ordering. The two terms in the Hamiltonian (13) account for the kinetic energy of the electrons and charging energy, respectively. Both scattering from a barrier in the constriction, and scattering from within a finite-size dot, create terms $\propto (\psi_R^\dagger \psi_L + h.c.)$ in the Hamiltonian, and in the corresponding action [10]. The backscattering terms lead to essentially non-perturbative corrections to the ground state energy E_0 . In the work of Matveev [5], the problem with a single scatterer in the constriction described by

$$\hat{H}_{bs} = v_F \left(r \psi_L^\dagger \psi_R + r^* \psi_R^\dagger \psi_L \right) \Big|_{x=0}, \quad (14)$$

was solved, and E_0 was found. According to [5], the regular Coulomb blockade oscillations in the energy are given by $E_{osc}(\mathcal{N}) \sim 2E_C |r|^2 \cos(2\pi\mathcal{N}) \ln [1/|r|^2 \cos^2(\pi\mathcal{N})]$ at $r \neq 0$.

Backscattering from inside the cavity ($\Delta \neq 0$) generates a random contribution to E_0 , superimposed on the regular oscillations. Once the reflection in the channel is accounted for non-perturbatively, the random backscattering from the walls of the dot can be studied by means of the lowest-order perturbation theory [10]:

$$\delta E_0 = i \frac{v_F^3}{\nu_{l0}} \int_{-\infty}^{\infty} dt \delta \mathcal{G}_{d0}(t) G_{LR}(-t; \mathcal{N}) \Big|_{\mathbf{r}=\mathbf{R}_c}, \quad (15)$$

where G_{LR} is a component of the electron Green function in a 1D infinite wire, described by the Hamiltonian $\hat{H} = \hat{H}_{1D} + \hat{H}_{bs}$ [see Eqs. (13), (14)],

$$G_{LR}(t) = \langle \psi_R^\dagger(t) \psi_L(0) + \psi_L^\dagger(t) \psi_R(0) \rangle \Big|_{x=0}. \quad (16)$$

Function G_{LR} describes the dynamics of backscattering of an individual electron from the dot. An electron wave in the channel carries charge and spin degrees of freedom,

and there are two time scales involved in the backscattering process. The shorter one, $t_\rho \sim \hbar/E_C$, corresponds to the reflection of the charge by the Coulomb barrier (resulting from the charging energy). The longer one, t_σ , describes the reflection of the spin component, and at finite r is of the order of $1/E_{osc}(\mathcal{N})$. At the intermediate time scale, $1/E_C < |t| < t_\sigma$, the charge and spin components of the backscattered electron are separated, and the Green function manifests a distinctively non-Fermi-liquid behavior,

$$G_{LR}(t; \mathcal{N}) = \frac{\cos \pi \mathcal{N}}{\pi v_F t} \sqrt{\frac{\gamma}{\pi^3} \frac{|t|}{t_\sigma(\mathcal{N})}} \ln \left(\frac{t_\sigma(\mathcal{N})}{|t|} \right). \quad (17)$$

Here $\gamma = e^{\mathbf{C}}$ with $\mathbf{C} = 0.577$ being the Euler constant. The electron backscattering process is completed at $t \sim t_\sigma$, and at larger time scales the system exhibits the standard Fermi-liquid behavior with $G_{LR}(t) \sim 1/t$. The result (17) can be obtained in the first-order perturbation theory in $\hat{H}_{bs}$, with the low-energy cut-off $E \sim E_{osc}(\mathcal{N})$.

Equations (2) and (15) enable us to relate capacitance fluctuations δC_{diff} to one-electron properties of the system described by the functions $\delta \mathcal{G}_{d0}$ and G_{LR} . The major contribution to the integral (15) is given by the interval $1/E_C < |t| < t_\sigma$, where approximation (17) is applicable. The correlation function of the capacitance fluctuations is then calculated with the help of Eqs. (7) and (8):

$$\begin{aligned} \langle \delta C_{diff}(\mathcal{N}, B_1) \delta C_{diff}(\mathcal{M}, B_2) \rangle &= \frac{16\gamma}{3\pi^2} C^2 \frac{\Delta}{E_C} |r|^2 \\ &\times \cos 2\pi \mathcal{N} \cos 2\pi \mathcal{M} \ln \left(\frac{1}{|r|^2 \cos^2 \pi \mathcal{N}} \right) \ln \left(\frac{1}{|r|^2 \cos^2 \pi \mathcal{M}} \right) \\ &\times \sum_{\pm} P \left(\frac{B_{\pm}^2}{B_c^2}; r_{max}^2 \right), \quad |r|^2 \equiv 1 - g, \end{aligned} \quad (18)$$

with $r_{max} \equiv \max\{|r \cos \pi \mathcal{N}|, |r \cos \pi \mathcal{M}|\}$ and $B_c \equiv (\Phi_0/S) \sqrt{2E_C/E_T}$. The variation of the correlation function with the magnetic field is described by the function $P(B^2/B_c^2; r_{max}^2)$. In the domain $B < B_c$,

$$P \left(\frac{B^2}{B_c^2}; r_{max}^2 \right) \simeq \left[\ln \frac{1}{r_{max}^2} \right]^3 - \left[\ln \max \left\{ 1; \frac{B^2}{r_{max}^2 B_c^2} \right\} \right]^3.$$

At magnetic fields larger than the correlation field B_c , the correlations rapidly decay, $P(B^2/B_c^2; r_{max}^2) \propto (B_c/B)^4$.

The perturbation theory Eq. (15) is applicable if the channel conductance is not too close to G_0 , *i.e.*, $|r|^2 \gg \Delta/E_C$. At a larger conductance, the amplitude of fluctuations saturates at $\delta C \sim C(\Delta/E_C) \ln^2(E_C/\Delta)$ [10].

Tunneling of arbitrary strength. The behavior of the capacitance fluctuations in the whole domain $0 \leq G_0 \leq G_q$ (with $G_q = e^2/\pi\hbar$) follows from Eqs. (9) and (18). At $G_0 \ll G_q$ the fluctuations increase with conductance, $\delta C_{diff}/C \sim (G_0/G_q) \sqrt{\Delta/E_C}$. For an almost open channel ($G_q - G_0 \ll G_q$) the characteristic amplitude of the capacitance fluctuations is $\delta C_{diff}/C \sim$

$[1 - (G_0/G_q)] \sqrt{\Delta/E_C}$, up to a $\ln^3$ factor. Thus the mesoscopic capacitance fluctuations reach maximum at a partial transmission of the channel connecting dot and lead. The parameter $\sqrt{\Delta/E_C}$ for typical semiconductor dots [13] is ~ 0.2 , so that the relative magnitude of these fluctuations is appreciable.

In conclusion, we found mesoscopic fluctuations of charge and differential capacitance of a partially opened quantum dot. The amplitude of fluctuations reaches maximum at the intermediate junction conductance, $G_0 \lesssim e^2/\pi\hbar$. We also found the correlation function of these fluctuations at different magnetic fields and gate voltages. In the experiments [6–8] no special attention was paid to the mesoscopic fluctuations. However, the parameters of quantum dots in [6–8] are in the right domain for studying mesoscopic fluctuations of the charge and of the differential capacitance. This can be accomplished by observing the Coulomb blockade with a magnetic field applied, which causes random variations of the charge.

The work at the University of Minnesota was supported by NSF Grants DMR-9423244 and DMR-9731756.

-
- [1] P. Lafarge, P. Joyez, D. Esteve, C. Urbina, M.H. Devoret, *Nature*, **365**, 422 (1993).
 - [2] L.I. Glazman and K.A. Matveev, *Sov. Phys. JETP* **71**, 1031 (1990), K.A. Matveev *ibid.* **72**, 892 (1991).
 - [3] G. Falci, G. Schön, G.T. Zimanyi, *Phys. Rev. Lett.*, **74**, 3257 (1995).
 - [4] K. Flensberg, *Phys. Rev. B* **48**, 11156 (1993); *Physica B* **203**, 432 (1994).
 - [5] K.A. Matveev, *Phys. Rev. B* **51**, 1743 (1995).
 - [6] F.R. Waugh, M.J. Berry, D.J. Mar, R.M. Westerwelt, K.L. Campman, and A.C. Gossard, *Phys. Rev. Lett.* **75**, 705 (1995).
 - [7] L.W. Molenkamp, K. Flensberg, M. Kemernik, *Phys. Rev. Lett.* **75**, 4282 (1995).
 - [8] D. Berman, N.V. Zhitenev, R.C. Ashoori, private communication.
 - [9] V.A. Gopar, P.A. Mello, and M. Büttiker, *Phys. Rev. Lett.* **77** 3005 (1996).
 - [10] I.L. Aleiner, L.I. Glazman, preprint cond-mat/9710195.
 - [11] B.L. Altshuler and A.G. Aronov, in *Electron-Electron Interaction in Disordered Conductors*, edited by A.L. Efros and M. Pollak (Elsevier, Amsterdam, 1985).
 - [12] In the elastic cotunneling process, [I.L. Aleiner, L.I. Glazman, *Phys. Rev. Lett.* **77**, 2057 (1996); S.M. Cronenwett, S.R. Patel, C.M. Marcus, K. Campman, and A.C. Gossard, *Phys. Rev. Lett.* **79**, 2312 (1997)] the correlation magnetic field is also determined by Eq. (10), with the same physics involved.
 - [13] L.P. Kouwenhoven, C.M. Marcus, P.L. McEuen, S. Tarucha, R.M. Westerwelt, N.S. Wingreen, in *Mesoscopic Electron Transport*, edited by L.L. Sohn, L.P. Kouwenhoven and G. Schön (Kluwer, Dordrecht, 1997).